



South Orange County Orthopaedics  
 26730 Crown Valley Parkway, Suite 200  
 Mission Viejo, CA 92691  
 www.SOCOrtho.com

**Roger C. Sohn, MD**  
 Sports Medicine/Arthroscopy, Disorders  
 of the Hand and Upper Extremity

## ACL RECONSTRUCTION PROTOCOL

(\* if a mensical repair is also performed, flexion should be limited to <90° until 6 weeks post-op)

<u>PHASE</u>	<u>GOAL</u>	<u>EXERCISE/METHODS</u>
<b>Phase 1</b> <b>Preoperative</b>	Decrease swelling Restore full terminal knee extension and flexion	Cold therapy with elevation (10-15 min) Extension: prone hangs or heel props (7-10 min) Flexion: wall slides, heel slides, active assist flexion
	Normalize gait	Weight bearing as tolerated with brace Treadmill: forward/backward (5-10 min)
	Normalize strength	Closed chain: leg press, total gym (10-50°), toe raises Open chain: Quad machine (90-30°), hamstring machine 4-way SLR. All exercises: 3-5 sets, 12-15 reps Electrical stimulation for VMO PRN
	Maintain aerobic fitness	Low-impact activities (bike, treadmill walking)
	Home program	Ice/elevation: 20 min on/60 min off Passive extension/flexion activities Lower extremity strengthening as indicated All activities 3-4 times/day or as indicated
<b>Phase 2</b> <b>Postoperative</b> <b>7-14 Days</b>	Control swelling Restore full terminal knee extension and flexion to 90°	Ace wrap with elevation-ankle pumps (20 min) Extension: heel props (7-10 min) Flexion: heel slides (7-10 min) *avoid prolonged lying with pillow under knee*
	Control swelling	Ace wrap with elevation-ankle pumps (20 min)  Treadmill: 5-10 min Calf, hamstring, hip flexor stretching (3 times, 30 sec hold) *emphasize heel to toe gait*

	Normalize gait	Weight bearing as tolerated with brace locked in full extension
	Increase strength	4-way SLR, quad sets, multi-angle isometrics (0-45-90°) Electrical muscle stimulation PRN (15 min) Partial squat (10-40°), toe raises with assistance PRN All exercises 3-5 sets, 12-15 reps Total Gym Level 1-3
	Home program	Ice/elevation: 20 min on/60 min off Passive extension/flexion activities Lower extremity strengthening as indicated All activities 3-4 times/day or as indicated *avoid prolonged lying with pillow under knee*
<b>Phase 3 Postoperative Weeks 2-5</b>	Control swelling	Ice PRN. Ace wrap PRN.
	Full passive knee extension and flexion (0-125 by 4 wks)( *if meniscus repaired flexion should be <90°)	Extension: prone hangs or heel props (7-10) Flexion: wall slides, heel slides, active assist flexion, bike (7-10 min)
	Normal ambulation	Treadmill: 5-10 min Increase grade with treadmill as tolerated Calf, hamstring, hip flexor stretching (3 times 30 sec hold)
	Pain-free strengthening	Closed chain: leg press (10-50), Total gym level (4-5) Push FROM without pain hamstring stretch, 4-way SLR, Toe raises, step ups, mini squats All exercises 3-5 sets, 12-15 reps *Implement within painfree ROM once normal gait and effusion control are achieved HS curls start at – 2 weeks for bone–patella–bone grafts and 4 weeks for HS grafts.
	Initiate proprioception exercises (pain-free)	*BAPS (start with ball 1 sitting and progress to standing - 3-4 min) *Double leg stand with rebounder *Single leg stance/grid exercises (lunges, reaches) with eyes open/eyes closed *Standing 4-way hip theratubing exercises *Lateral treadmill walking both directions *Implement once patient is able to ambulate normally without pain (3-5 min)
<b>Phase 4 Postoperative Week 6-12</b>	Criteria for progression to Phase 5 No effusion, painless full ROM, minimal crepitus. Score greater than 80 On Lysholm with no locking or instability	
	Improve aerobic level of fitness	Painfree low impact activity (bike, treadmill walking, stepper on own) 3-4 times per week, 20-30 minutes
	Improve proprioception	Trampoline hopping bilaterally progressing To jogging, then single leg hopping (10 min) (When Quadricep strength approaches 65%). Fitter with ski poles progressing to no ski poles (7-10 minutes) BAPS – single leg

Body blade Proprioceptive Star Exercises  
 [8-12 weeks]  
 \*Week 10 – Light double leg hopping  
 (multi-directional), lateral shuffles  
 \*Only if all previous activities were  
 completed without pain or increased  
 swelling

Improve Strength

Total gym level 8-10 without pain, leg press  
 (10-50), toe raises, step ups  
 Hamstring machine, 4-way SLR  
 All exercises 3-5 sets, 12-15 reps  
 Mini squats, Partial lunges,  
 single-leg toe raise

**Phase 6  
 Postoperative  
 Week 13-24**

Criteria for progression to Phase 6  
 no effusion, painless full ROM,  
 minimal crepitus. Score greater than  
 90 on Lysholm with no locking or instability  
 Ratio of 80% involved to uninjured on isokinetic test at 6 months post-op  
 Ratio of 80% on one-legged jump

Improve aerobic level of fitness  
 (Usually 3-4 months)

Painfree low impact activity (bike, treadmill  
 Walking, stepper on own) 3-4 times per  
 Week, 20-30 minutes

Improve strength

Closed chain: leg press (10-50°),  
 Toe raises. Total gym level 8-10 without  
 Pain.  
 Open chain: Quad machine (90-30°),  
 Hamstring machine  
 All exercises: 3 sets 15-20 each

Implement running program if indicated

Walk/jog cycles  
 3-5 min warm-up (walk)  
 \*10-30 sec jog/60 sec walk for 10-20 min  
 3-5 min cool-down (walk)  
 \*increase jog time and decrease walk time  
 based upon patient response.

Implement sport specific activity

\*Progression program of strength training  
 And agility/speed drills  
 \*Program should be based upon specific  
 patient needs.